

Section 1.9 (Through Theorem 10) The Matrix of a Linear Transformation

Identity Matrix I_n is an $n \times n$ matrix with 1's on the main left to right diagonal and 0's elsewhere. The i th column of I_n is labeled \mathbf{e}_i .

EXAMPLE:

$$I_3 = \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Note that

$$I_3 \mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$= \text{---} \begin{bmatrix} \\ \\ \end{bmatrix} + \text{---} \begin{bmatrix} \\ \\ \end{bmatrix} + \text{---} \begin{bmatrix} \\ \\ \end{bmatrix} = \text{---}.$$

In general, for \mathbf{x} in \mathbf{R}^n ,

$$I_n \mathbf{x} = \text{---}$$

From Section 1.8, if $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ is a linear transformation, then $T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v})$.

Generalized Result:

$$T(c_1\mathbf{v}_1 + \cdots + c_p\mathbf{v}_p) = c_1T(\mathbf{v}_1) + \cdots + c_pT(\mathbf{v}_p).$$

EXAMPLE: The columns of $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ are $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Suppose T is a linear transformation from \mathbf{R}^2 to \mathbf{R}^3 where

$$T(\mathbf{e}_1) = \begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix} \text{ and } T(\mathbf{e}_2) = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}.$$

Compute $T(\mathbf{x})$ for any $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$.

Solution: A vector \mathbf{x} in \mathbf{R}^2 can be written as

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \text{---} \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \text{---} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \text{---} \mathbf{e}_1 + \text{---} \mathbf{e}_2$$

Then

$$T(\mathbf{x}) = T(x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2) = \text{---} T(\mathbf{e}_1) + \text{---} T(\mathbf{e}_2)$$

$$= \text{---} \begin{bmatrix} 2 \\ -3 \\ 4 \end{bmatrix} + \text{---} \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}.$$

Note that

$$T(\mathbf{x}) = \begin{bmatrix} \\ \\ \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}.$$

So

$$T(\mathbf{x}) = \begin{bmatrix} T(\mathbf{e}_1) & T(\mathbf{e}_2) \end{bmatrix} \mathbf{x} = A\mathbf{x}$$

To get A , replace the identity matrix $\begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 \end{bmatrix}$ with $\begin{bmatrix} T(\mathbf{e}_1) & T(\mathbf{e}_2) \end{bmatrix}$.

Theorem 10

Let $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation. Then there exists a unique matrix A such that

$$T(\mathbf{x}) = A\mathbf{x} \text{ for all } \mathbf{x} \text{ in } \mathbf{R}^n.$$

In fact, A is the $m \times n$ matrix whose j th column is the vector $T(\mathbf{e}_j)$, where \mathbf{e}_j is the j th column of the identity matrix in \mathbf{R}^n .

$$A = [T(\mathbf{e}_1) \quad T(\mathbf{e}_2) \quad \cdots \quad T(\mathbf{e}_n)]$$

↑

standard matrix for the linear transformation T


EXAMPLE:
$$\begin{bmatrix} ? & ? \\ ? & ? \\ ? & ? \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 2x_2 \\ 4x_1 \\ 3x_1 + 2x_2 \end{bmatrix}$$

Solution:

$$\begin{bmatrix} ? & ? \\ ? & ? \\ ? & ? \end{bmatrix} = \text{standard matrix of the linear transformation } T$$

$$\begin{bmatrix} ? & ? \\ ? & ? \\ ? & ? \end{bmatrix} = \begin{bmatrix} T(\mathbf{e}_1) & T(\mathbf{e}_2) \end{bmatrix} =$$

EXAMPLE: Find the standard matrix of the linear transformation $T : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ which rotates a point about the origin through an angle of $\frac{\pi}{4}$ radians (counterclockwise).



$$T(\mathbf{e}_1) = \begin{bmatrix} \\ \end{bmatrix}$$

$$T(\mathbf{e}_2) = \begin{bmatrix} \\ \end{bmatrix}$$

$$\Rightarrow A = \begin{bmatrix} & \\ & \end{bmatrix}$$